arXiv:1409.7152v4 [math.RA] 23 Mar 2015

The Drinfel'd Double versus the Heisenberg Double for Hom-Hopf Algebras

Daowei Lu, Shuanhong Wang *

Department of Mathematics, Southeast University

Nanjing, Jiangsu 210096, P. R. of China

Abstract. Let (A, α) be a finite-dimensional Hom-Hopf algebra. In this paper we mainly construct the Drinfel'd double $D(A) = (A^{op} \bowtie A^*, \alpha \otimes (\alpha^{-1})^*)$ in the setting of Hom-Hopf algebras by two ways, one of which generalizes Majid's bicrossproduct for Hopf algebras (see [7]) and another one is to introduce the notion of dual pairs of of Hom-Hopf algebras. Then we study the relation between the Drinfel'd double $D(A)$ and Heisenberg double $H(A) = A\#A^*$, generalizing the main result in [5]. Especially, the examples given in the paper are not obtained from the usual Hopf algebras.

Keywords: Hom-Hopf algebra; Drinfel'd double; Majid's bicrossproduct; Heisenberg double.

Mathematics Subject Classification: 16W30.

0. INTRODUCTION

As well-known, in physics, the Yang-Baxter equation (or star-triangle relation) is a consistency equation which was first introduced in the field of statistical mechanics, and it takes its name from independent work of C. N. Yang in 1968, and R. J. Baxter in 1971. It depends on the idea that in some scattering situations, particles may preserve their momentum while changing their quantum internal states. It states that a matrix R, acting on two out of three objects, satisfies

$$(R \otimes id)(id \otimes R)(R \otimes id) = (id \otimes R)(R \otimes id)(id \otimes R).$$

*Corresponding author: Shuanhong Wang, shuanhwang@seu.edu.cn, Department of Mathematics, Southeast University, Nanjing, Jiangsu 210096, P. R. of China.

In one dimensional quantum systems, R is the scattering matrix and if it satisfies the Yang-Baxter equation then the system is integrable. The Yang-Baxter equation also shows up when discussing knot theory and the braid groups where R corresponds to swapping two strands. Since one can swap three strands two different ways, the Yang-Baxter equation enforces that both paths are the same.

Braided monoidal categories give rise to solutions to the quantum Yang-Baxter equation (QYBE). The classical braided monoidal categories come from the representations of quasitriangular Hopf algebras. Quasitriangular Hopf algebras have been very widely studied and one of the most important examples of a quasitriangular Hopf algebra is the Drinfel'd double (see [2]). As such, they are interesting to different research communities in mathematical physics (see [7] for example). Although the Drinfel'd double for finite-dimensional Hopf algebras provide examples of such solutions to the QYBE, these are rather trivial. The wish to obtain more interesting solutions to the QYBE provides a strong motivation to find new examples of Drinfel'd double in the setting of Hom-Hopf algebras, whose representations are a braided monoidal category.

The aim of this article is to construct new examples of braided monoidal categories in the setting of Hom-Hopf algebras. This is achieved by generalizing an existing the Drinfel'd double (see [2]). We will generalize this construction to the so-called Hom-Hopf algebras, which are Hopf algebras in the Hom-category of a monoidal category (see [1]). We find a suitable generalization of the notions of a Majid's bicrossproduct and dual pairs for this setting and obtain a Drinfel'd double of a Hom-Hopf algebra whose representation is a braided monoidal category. Furthermore, we establish the relation between Drinfel'd double and Heisenberg double in the setting of Hom-Hopf algebras, generalizing the results in [5]. Especially, the examples obtained in the paper are not obtained from the usual Hopf algebras.

The article is organized as follows.

In Section 1, we will recall the definitions and results of Hom-Hopf algebras, such as Hom-algebas, Hom-coalgebras, Hom-modules, Hom-comodules and the Hom-smash products.

Let (A, α_A) and (H, α_H) be finite-dimensional Hom-Hopf algebras. In Section 2, we will introduce the notion of a bicrossproduct $(A \ltimes H, \alpha_A \otimes \alpha_H)$, and give the conditions for $(A \ltimes H, \alpha_A \otimes \alpha_H)$ to form a Hom-Hopf algebra, generalizing the Majid's bicrossproduct defined in [6] (see Theorem 2.6). And examples of bicrossproduct Hom-Hopf algebras

are constructed. Then we will show how to get the Drinfel'd double by our new theory. Furthermore solutions of the QHYBE are obtained.

In Section 3, we will construct the Drinfel'd double associated to a dual pair of Hom-Hopf algebras (see Theorem 3.4), which is a generalization of the Drinfel'd double in the finite dimensional case. In section 4, we will establish the relation between Drinfel'd double and Heisenberg double (see Theorem 4.5). Examples are given to illustrate our theory.

1. PRELIMINARIES

Throughout this article, all the vector spaces, tensor product and homomorphisms are over a fixed field k. For a coalgebra C, we will use the Heyneman-Sweedler's notation $\Delta(c) = c_1 \otimes c_2$, for any $c \in C$ (summation omitted).

In this section, we will recall the definitions in [8] on the Hom-Hopf algebras, Hom-modules and Hom-comodules.

A unital Hom-associative algebra is a triple (A, μ, α) where $\alpha : A \longrightarrow A$ and $\mu : A \otimes A \longrightarrow A$ are linear maps, with notation $\mu(a \otimes b) = ab$ such that for any $a, b, c \in A$,

$$\begin{aligned} &\alpha(ab) = \alpha(a)\alpha(b),\ \alpha(1_A) = 1_A, \\ &1_A a = \alpha(a) = a1_A,\ \alpha(a)(bc) = (ab)\alpha(c). \end{aligned}$$

A linear map $f : (A, \mu_A, \alpha_A) \longrightarrow (B, \mu_B, \alpha_B)$ is called a morphism of Hom-associative algebra if $\alpha_B \circ f = f \circ \alpha_A$, $f(1_A) = 1_B$ and $f \circ \mu_A = \mu_B \circ (f \otimes f)$.

A counital Hom-coassociative coalgebra is a triple $(C, \Delta, \varepsilon, \alpha)$ where $\alpha : C \longrightarrow C$, $\varepsilon : C \longrightarrow k$, and $\Delta : C \longrightarrow C \otimes C$ are linear maps such that

$$\begin{aligned} &\varepsilon \circ \alpha = \varepsilon,\ (\alpha \otimes \alpha) \circ \Delta = \Delta \circ \alpha, \\ &(\varepsilon \otimes id) \circ \Delta = \alpha = (id \otimes \varepsilon) \circ \Delta, \\ &(\Delta \otimes \alpha) \circ \Delta = (\alpha \otimes \Delta) \circ \Delta. \end{aligned}$$

A linear map $f : (C, \Delta_C, \alpha_C) \longrightarrow (D, \Delta_D, \alpha_D)$ is called a morphism of Hom-coassociative coalgebra if $\alpha_D \circ f = f \circ \alpha_C$, $\varepsilon_D \circ f = \varepsilon_C$ and $\Delta_D \circ f = (f \otimes f) \circ \Delta_C$.

In what follows, we will always assume all Hom-algebras are unital and Hom-coalgebras are counital.

A Hom-bialgebra is a quadruple (H, μ, Δ, α), where (H, μ, α) is a Hom-associative algebra and (H, Δ, α) is a Hom-coassociative coalgebra such that Δ and ε are morphisms of Hom-associative algebra.

A Hom-Hopf algebra (H, μ, Δ, α) is a Hom-bialgebra H with a linear map $S : H \longrightarrow H$(called antipode) such that

$$S \circ \alpha = \alpha \circ S,$$
$$S(h_1)h_2 = h_1 S(h_2) = \varepsilon(h)1,$$

for any $h \in H$. For S we have the following properties:

$$S(h)_1 \otimes S(h)_2 = S(h_2) \otimes S(h_1),$$
$$S(gh) = S(h)S(g),\ \varepsilon \circ S = \varepsilon.$$

For any Hopf algebra H and any Hopf algebra endomorphism α of H, there exists a Hom-Hopf algebra $H_\alpha = (H, \alpha \circ \mu, 1_H, \Delta \circ \alpha, \varepsilon, S, \alpha)$.

Let (A, α_A) be a Hom-associative algebra, M a linear space and $\alpha_M : M \longrightarrow M$ a linear map. A left A-module structure on (M, α_M) consists of a linear map $A \otimes M \longrightarrow M$, $a \otimes m \mapsto a \cdot m$, such that

$$1_A \cdot m = \alpha_M(m),$$
$$\alpha_M(a \cdot m) = \alpha_A(a) \cdot \alpha_M(m),$$
$$\alpha_A(a) \cdot (b \cdot m) = (ab) \cdot \alpha_M(m),$$

for any $a, b \in A$ and $m \in M$.

Similarly we can define the right (A, α)-modules. Let (M, μ) and (N, ν) be two left (A, α)-modules, then a linear map $f : M \longrightarrow N$ is a called left A-module map if $f(am) = af(m)$ for any $a \in A$, $m \in M$ and $f \circ \mu = \nu \circ f$.

Let (C, α_C) be a Hom-coassociative coalgebra, M a linear space and $\alpha_M : M \longrightarrow M$ a linear map. A right C-comodule structure on (M, α_M) consists a linear map $\rho : M \longrightarrow M \otimes C$ such that

$$(id \otimes \varepsilon_C) \circ \rho = \alpha_M,$$
$$(\alpha_M \otimes \alpha_C) \circ \rho = \rho \circ \alpha_M,$$
$$(\rho \otimes \alpha_C) \circ \rho = (\alpha_M \otimes \Delta) \circ \rho.$$

Let (M, μ) and (N, ν) be two right (C, γ)-comodules, then a linear map $g : M \longrightarrow N$ is a called right C-comodule map if $g \circ \mu = \nu \circ g$ and $\rho_N \circ g = (g \otimes id) \circ \rho_M$.

Let $(H, \mu_H, \Delta_H, \alpha_H)$ be a Hom-bialgebra. A Hom-associative algebra (A, μ_A, α_A) is called a left H-module Hom-algebra if (A, α_A) is a left H-module, with the action $H \otimes A \longrightarrow A,\ h \otimes a \mapsto h \cdot a$, such that

$$\alpha_H^2(h) \cdot (ab) = (h_1 \cdot a)(h_2 \cdot b),$$

$$h \cdot 1_A = \varepsilon(h)1_A,$$

for any $h \in H$ and $a, b \in A$.

When A is a left H-module Hom-algebra, in [8] the Hom-smash product $A\#H$ is defined as follows:

$$(a\#h)(b\#k) = a(\alpha_H^{-2}(h_1) \cdot \alpha_A^{-1}(b))\#\alpha_H^{-1}(h_2)k,$$

for any $a, b \in A$ and $h, k \in H$.

Recall from [12] that a Hom-bialgebra (H, α_H) is quasitriangular if there exists an element $R \in H \otimes H$ satisfying:

1. $\Delta^{op}(x)R = R\Delta(x)$ for any $x \in H$,
2. $(\Delta \otimes \alpha_H)R = R^{13}R^{23}$,
3. $(\alpha_H \otimes \Delta)R = R^{13}R^{12}$.

2. MAJID'S BICROSSPRODUCTS FOR HOM-HOPF ALGEBRAS

In this section, we will construct the bicrossproduct on Hom-bialgebras, which generalize the Majid's bicrossproducts in the usual Hopf algebras. Then we can get Drinfel'd double for Hom-Hopf algebras in the setting of the Majid's bicrossproducts.

In [4] L. Liu and B. Shen defined the smash coproduct on monoidal Hom-Hopf algebra. Now we need to construct the smash coproduct on Hom-Hopf algebra.

Definition 2.1. Let (H, α_H) be a Hom-bialgebra. A Hom-coassociative coalgebra (C, α_C) is called a right *H-comodule Hom-coalgebra* if (C, α_C) a right H-comodule, with the comodule structure map $\rho : C \longrightarrow C \otimes H$, $c \mapsto c_{(0)} \otimes c_{(1)}$, such that the following conditions hold

$$\varepsilon_C(c_{(0)})c_{(1)} = \varepsilon_C(c)1_H,$$
$$c_{(0)1} \otimes c_{(0)2} \otimes \alpha_H^2(c_{(1)}) = c_{1(0)} \otimes c_{2(0)} \otimes c_{1(1)}c_{2(1)},$$

for any $c \in C$.

Example 2.2. (1) Let H be a bialgebra and C a right H-comodule coalgebra in the usual sense. The coaction is denoted by $C \longrightarrow C \otimes H$, $c \mapsto c_{[0]} \otimes c_{[1]}$. Let α_H be a bialgebra endomorphism of H, and α_C a coalgebra endomorphism of C such that

$$\alpha_C(c)_{[0]} \otimes \alpha_C(c)_{[1]} = \alpha_C(c_{[0]}) \otimes \alpha_H(c_{[1]}),$$

for any $c \in C$. Consider the Hom-bialgebra $H_{\alpha_H} = (H, \alpha_H \circ \mu_H, \Delta_H \circ \alpha_H, \alpha_H)$ and the Hom-coassociative coalgebra $C_{\alpha_C} = (C, \Delta_C \circ \alpha_C, \alpha_C)$. Then C_{α_C} is a right H_{α_H}-comodule Hom-coalgebra with the coaction $C_{\alpha_C} \longrightarrow C_{\alpha_C} \otimes H_{\alpha_H}$, $c \mapsto c_{(0)} \otimes c_{(1)} = \alpha_C(c_{[0]}) \otimes \alpha_H(c_{[1]})$.

(2) Let (H, α) be any Hom-Hopf algebra. Then (H^{op}, α) is a right (H, α)-comodule Hom-coalgebra with the right coaction

$$\rho(h) = \alpha^{-1}(h_{12}) \otimes S(\alpha^{-2}(h_{11}))\alpha^{-1}(h_2),$$

for any $h \in H$. The fact that (H^{op}, α) is a right (H, α)-comodule is obvious, and for $h \in H$,

$$\begin{aligned}
&h_{(0)1} \otimes h_{(0)2} \otimes \alpha^2(h_1)\\
=&\alpha^{-1}(h_{121}) \otimes \alpha^{-1}(h_{122}) \otimes S(h_{11})\alpha(h_2)\\
=&h_{12} \otimes \alpha^{-1}(h_{212}) \otimes [S(\alpha^{-1}(h_{11}))(\alpha^{-4}(h_{2111})S(\alpha^{-4}(h_{2112})))]h_{22}\\
=&\alpha^{-1}(h_{112}) \otimes \alpha^{-1}(h_{212}) \otimes [S(\alpha^{-2}(h_{111}))(\alpha^{-2}(h_{12})S(\alpha^{-3}(h_{211})))]h_{22}\\
=&\alpha^{-1}(h_{112}) \otimes \alpha^{-1}(h_{212}) \otimes [S(\alpha^{-2}(h_{111}))\alpha^{-1}(h_{12})][S(\alpha^{-2}(h_{211}))\alpha^{-1}(h_{22})]\\
=&h_{1(0)} \otimes h_{2(0)} \otimes h_{1(1)}h_{2(1)}.
\end{aligned}$$

Then (H^{op}, α) is a right (H, α)-comodule Hom-coalgebra.

(3) Let $A_x^1 = span\{1, x\}$ over a fixed field k with $chark \neq 2$. Then define β as a k-linear automorphism of A_x^1 by

$$\beta(1_{A_x^1}) = 1_{A_x^1}, \quad \beta(x) = -x.$$

Define the multiplication on A_x^1 by

$$1_{A_x^1} 1_{A_x^1} = 1_{A_x^1},\ 1_{A_x^1} x = -x,\ x^2 = 0,$$

then it is not hard to check that (A_x^1, β) is a Hom-associative algebra.

For A_x^1, define the coalgebra structure and antipode by

$$\Delta(1_{A_x^1}) = 1_{A_x^1} \otimes 1_{A_x^1},\ \Delta(x) = (-x) \otimes 1 + 1 \otimes (-x),$$

$$\varepsilon(1_{A_x^1}) = 1,\ \varepsilon(x) = 0, \quad S_{A_x^1}(1_{A_x^1}) = 1_{A_x^1},\ S_{A_x^1}(x) = -x,$$

then (A_x^1, β) is a Hom-Hopf algebra.

Let $H = span\{1, g | g^2 = 1\}$ be the group algebra. Obviously (H, id) is a Hom-Hopf algebra. Define the left action of H on A_x^1 $\cdot : H \otimes A_x^1 \longrightarrow A_x^1$ by

$$1_H \cdot 1_{A_x^1} = 1_{A_x^1},\ 1_H \cdot x = -x,\ g \cdot 1_{A_x^1} = 1_{A_x^1},\ g \cdot x = x.$$

It is easy to check that (A_x^1, β) is a left (H, id)-module algebra.

Define the right coaction of A_x^1 on H $\rho : H \longrightarrow H \otimes A_x^1$ by

$$\rho(1_H) = 1_H \otimes 1_{A_x^1},\ \rho(g) = g \otimes 1_{A_x^1},$$

then (H, id) is a right (A_x^1, β)-comodule coalgebra.

Definition 2.3. Let (C, Δ_C, α_C) and (D, Δ_D, α_D) be Hom-coassociative coalgebras. A linear map $\Phi : C \otimes D \longrightarrow D \otimes C$ is called a *Hom-cotwisting map* between C and D if the following conditions hold:

$$(\Delta_D \otimes \alpha_C) \circ \Phi = (id_D \otimes \Phi) \circ (\Phi \otimes id_D) \circ (\alpha_C \otimes \Delta_D), \tag{2. 1}$$

$$(\alpha_D \otimes \Delta_C) \circ \Phi = (\Phi \otimes id_C) \circ (id_C \otimes \Phi) \circ (\Delta_C \otimes \alpha_D), \tag{2. 2}$$

$$(\alpha_D \otimes \alpha_C) \circ \Phi = \Phi \circ (\alpha_C \otimes \alpha_D), \tag{2. 3}$$

$$(\varepsilon_C \otimes id) \circ \Phi = id \otimes \varepsilon_C,\ (id \otimes \varepsilon_D) \circ \Phi = \varepsilon_D \otimes id. \tag{2. 4}$$

If we denote the element $\Phi(c \otimes d)$ in $D \otimes C$ by $\Phi(c \otimes d) = d^\Phi \otimes c^\Phi = d^\phi \otimes c^\phi$ for any $c \in C, d \in D$, the above identities can be rewritten as

$$\begin{cases} (d^\Phi)_1 \otimes (d^\Phi)_2 \otimes \alpha_C(c^\Phi) = (d_1)^\Phi \otimes (d_2)^\phi \otimes \alpha_C(c)^{\Phi\phi}, \\ \alpha_D(d^\Phi) \otimes (c^\Phi)_1 \otimes (c^\Phi)_2 = \alpha_D(d)^{\Phi\phi} \otimes (c_1)^\phi \otimes (c_2)^\Phi, \\ \alpha_D(d^\Phi) \otimes \alpha_C(c^\Phi) = \alpha_D(d)^\Phi \otimes \alpha_C(c)^\Phi, \\ \varepsilon_C(c^\Phi)d^\Phi = \varepsilon_C(c)d,\ \varepsilon_D(d^\Phi)c^\Phi = \varepsilon_D(d)c. \end{cases}$$

Proposition 2.4. Let (C, Δ_C, α_C) and (D, Δ_D, α_D) be Hom-coassociative coalgebras and $\Phi : C \otimes D \longrightarrow D \otimes C$ a Hom-cotwisting map. Define $\Delta : C \otimes D \longrightarrow (C \otimes D) \otimes (C \otimes D)$ by

$$\Delta(c \otimes d) = c_1 \otimes d_1^\Phi \otimes c_2^\Phi \otimes d_2,$$

and $\varepsilon : C \otimes D \longrightarrow k$ by

$$\varepsilon(c \otimes d) = \varepsilon_C(c)\varepsilon_D(d),$$

for any $c \in C, d \in D$. Then $(C \otimes D, \Delta, \alpha_C \otimes \alpha_D)$ is a Hom-coassociative coalgebra.

Proof. The counit is straightforward to verify. For any $c \in C, d \in D$, firstly

$$\begin{aligned} (\Delta \circ (\alpha_C \otimes \alpha_D))(c \otimes d) &= \Delta(\alpha_C(c) \otimes \alpha_D(d)) \\ &= \alpha_C(c_1) \otimes \alpha_D(d_1)^\Phi \otimes \alpha_C(c_2)^\Phi \otimes \alpha_D(d_2) \\ &= \alpha_C(c_1) \otimes \alpha_D(d_1^\Phi) \otimes \alpha_C(c_2^\Phi) \otimes \alpha_D(d_2) \\ &= ((\alpha_C \otimes \alpha_D) \otimes (\alpha_C \otimes \alpha_D)) \circ \Delta(c \otimes d). \end{aligned}$$

Then

$$\begin{aligned}
& ((\alpha_C \otimes \alpha_D) \otimes \Delta) \circ \Delta(c \otimes d) \\
= \quad & \alpha_C(c_1) \otimes \alpha_D(d_1^{\Phi}) \otimes \Delta(c_2^{\Phi} \otimes d_2) \\
= \quad & \alpha_C(c_1) \otimes \alpha_D(d_1^{\Phi}) \otimes (c_2^{\Phi})_1 \otimes (d_{21})^{\phi} \otimes ((c_2^{\Phi})_2)^{\phi} \otimes d_{22} \\
\overset{(2.1)}{=} \quad & \alpha_C(c_1) \otimes \alpha_D(d_1)^{\Phi\phi'} \otimes (c_{21})^{\phi'} \otimes (d_{21})^{\phi} \otimes (c_{22})^{\Phi\phi} \otimes d_{22} \\
= \quad & c_{11} \otimes (d_{11})^{\Phi\phi'} \otimes (c_{12})^{\phi'} \otimes (d_{12})^{\phi} \otimes \alpha_C(c_2)^{\Phi\phi} \otimes \alpha_D(d_2) \\
\overset{(2.2)}{=} \quad & c_{11} \otimes ((d_1^{\Phi})_1)^{\phi} \otimes (c_{12})^{\phi} \otimes (d_1^{\Phi})_2 \otimes \alpha_C(c_2^{\Phi}) \otimes \alpha_D(d_2) \\
= \quad & (\Delta \otimes (\alpha_C \otimes \alpha_D)) \circ \Delta(c \otimes d).
\end{aligned}$$

By definition, this finishes the proof. ■

Example 2.5. (1) Let (H, α_H) be a Hom-bialgebra and (C, α_C) a right H-comodule Hom-coalgbra with the coaction $C \longrightarrow C \otimes H,\ c \mapsto c_{(0)} \otimes c_{(1)}$. Define the linear map $\Phi_1 : H \otimes C \longrightarrow C \otimes H$ by

$$\Phi_1(h \otimes c) = \alpha_C^{-1}(c_{(0)}) \otimes \alpha_H^{-1}(h)\alpha_C^{-2}(c_{(1)}),$$

for any $h \otimes H$ and $c \in C$. Then Φ_1 is a Hom-cotwisting map between H and C.

(2) Let (H, α) be a Hom-bialgebra and ρ the coaction of H on H^{op} in Example 2.2 (2), we have a Hom-cotwisting map $\Phi_2 : H \otimes H^{op} \longrightarrow H^{op} \otimes H$ given by

$$\Phi_2(k \otimes h) = \alpha^{-2}(h_{12}) \otimes \alpha^{-1}(k)[S(\alpha^{-4}(h_{11}))\alpha^{-3}(h_2)],$$

for any $h,\ k \in H$.

(3) With the notations in Example 2.2 (3), we have a Hom-cotwisting map $\Phi_3 : A_x^1 \otimes H \longrightarrow H \otimes A_x^1$ given by

$$\begin{aligned}
&\Phi_3(1_{A_x^1} \otimes 1_H) = 1_H \otimes 1_{A_x^1}, \quad \Phi_3(x \otimes 1_H) = 1_{A_x^1} \otimes -x, \\
&\Phi_3(1_{A_x^1} \otimes g) = g \otimes 1_{A_x^1}, \qquad \Phi_3(x \otimes g) = g \otimes -x.
\end{aligned}$$

Theorem 2.6. Let (A, S_A, α_A) and (H, S_H, α_H) be Hom-Hopf algebras. Assume that A is a left H-module Hom-algebra with the action $H \otimes A \longrightarrow A,\ h \otimes a \mapsto h \cdot a$ and H is a right A-comodule Hom-coalgebra with the coaction $H \longrightarrow H \otimes A,\ h \mapsto h_{(0)} \otimes h_{(1)}$. If the following conditions are satisfied:

(1) For any $h \in H, b \in A$,

$$\Delta(h \cdot b) = \alpha_H^{-1}(h_{1(0)}) \cdot b_1 \otimes \alpha_A^{-1}(h_{1(1)})(\alpha_H^{-1}(h_2) \cdot \alpha_A^{-1}(b_2)),$$

$$\varepsilon_A(h \cdot b) = \varepsilon_A(b)\varepsilon_H(h),$$

(2) For any $h, g \in H$,

$$(hg)_{(0)} \otimes (hg)_{(1)} = \alpha_H^{-1}(h_{1(0)})g_{(0)} \otimes \alpha_A^{-1}(h_{1(1)})(\alpha_H^{-1}(h_2) \cdot \alpha_A^{-1}(g_{(1)})),$$

(3) For any $h \in H, b \in A$,

$$h_{2(0)} \otimes (h_1 \cdot b)h_{2(1)} = h_{1(0)} \otimes h_{1(1)}(h_2 \cdot b),$$

then $(A \otimes H, \alpha_A \otimes \alpha_H)$ is a Hom-Hopf algebra under the Hom-smash product and Hom-smash coproduct, that is,

$$(a \otimes h)(b \otimes g) = a(\alpha_H^{-2}(h_1) \cdot \alpha_A^{-1}(b)) \otimes \alpha_H^{-1}(h_2)g,$$

and

$$\Delta(a \otimes h) = a_1 \otimes \alpha_H^{-1}(h_{1(0)}) \otimes \alpha_A^{-1}(a_2)\alpha_A^{-2}(h_{1(1)}) \otimes h_2,$$

with the antipode $S : A \otimes H \longrightarrow A \otimes H$ given by

$$S(a \otimes h) = (1 \otimes S_H(\alpha_H^{-2}(h_{(0)})))(S_A(\alpha_A^{-2}(a)\alpha_A^{-3}(h_{(1)})) \otimes 1),$$

for any $h, g \in H$, $a, b \in A$.

We will call this Hom-Hopf algebra Hom-bicrossproduct, i.e, *Majid's bicrossproduct* for Hom-Hopf algebras, and denote it by $(A \ltimes H, \alpha_A \otimes \alpha_H)$.

Proof. Firstly we need to prove Δ and ε is a morphism of Hom-associative algebra. Indeed for any $a, b \in A$ and $g, h \in H$,

$$\begin{aligned}
&\Delta((a \otimes h)(b \otimes g)) \\
=&\Delta(a(\alpha_H^{-2}(h_1) \cdot \alpha_A^{-1}(b)) \otimes \alpha_H^{-1}(h_2)g) \\
=&a_1(\alpha_H^{-2}(h_1) \cdot \alpha_A^{-1}(b)_1) \otimes \alpha_H^{-1}((\alpha_H^{-1}(h_{21})g_1)_{(0)}) \\
&\otimes [\alpha_A^{-1}(a_2)\alpha_A^{-1}((\alpha_H^{-2}(h_1) \cdot \alpha_A^{-1}(b))_2)]\alpha_A^{-2}((\alpha_H^{-1}(h_{21})g_1)_{(1)}) \otimes \alpha_H^{-1}(h_{22})g_2 \\
=&a_1(\alpha_H^{-3}(h_{11(0)}) \cdot \alpha_A^{-1}(b_1)) \otimes \alpha_H^{-3}(h_{211(0)})\alpha_H^{-1}(g_{1(0)}) \\
&\otimes \{\alpha_A^{-1}(a_2)[\alpha_A^{-4}(h_{11(1)})(\alpha_H^{-4}(h_{12}) \cdot \alpha_A^{-3}(b_2))]\} \\
&\{\alpha_A^{-4}(h_{211(1)})(\alpha_H^{-4}(h_{212}) \cdot \alpha_A^{-3}(g_{1(1)}))\} \otimes \alpha_H^{-1}(h_{22})g_2 \\
=&a_1(\alpha_H^{-3}(h_{11(0)}) \cdot \alpha_A^{-1}(b_1)) \otimes \alpha_H^{-3}(h_{211(0)})\alpha_H^{-1}(g_{1(0)}) \\
&\otimes \{\alpha_A^{-1}(a_2)\alpha_A^{-3}(h_{11(1)})\}\{[(\alpha_H^{-3}(h_{12}) \cdot \alpha_A^{-2}(b_2))] \\
&[\alpha_A^{-5}(h_{211(1)})(\alpha_H^{-5}(h_{212}) \cdot \alpha_A^{-4}(g_{1(1)}))]\} \otimes \alpha_H^{-1}(h_{22})g_2
\end{aligned}$$

$$
\begin{aligned}
=&a_1(\alpha_H^{-3}(h_{11(0)})\cdot\alpha_A^{-1}(b_1))\otimes\alpha_H^{-3}(h_{211(0)})\alpha_H^{-1}(g_{1(0)})\otimes\{\alpha_A^{-1}(a_2)\alpha_A^{-3}(h_{11(1)})\}\\
&\{[(\alpha_H^{-4}(h_{12})\cdot\alpha_A^{-3}(b_2))\alpha_A^{-5}(h_{211(1)})](\alpha_H^{-4}(h_{212})\cdot\alpha_A^{-3}(g_{1(1)}))\}\otimes\alpha_H^{-1}(h_{22})g_2\\
=&a_1(\alpha_H^{-2}(h_{1(0)})\cdot\alpha_A^{-1}(b_1))\otimes\alpha_H^{-4}(h_{2112(0)})\alpha_H^{-1}(g_{1(0)})\otimes\{\alpha_A^{-1}(a_2)\alpha_A^{-2}(h_{1(1)})\}\\
&\{[(\alpha_H^{-6}(h_{2111})\cdot\alpha_A^{-3}(b_2))\alpha_A^{-6}(h_{2112(1)})](\alpha_H^{-4}(h_{212})\cdot\alpha_A^{-3}(g_{1(1)}))\}\otimes\alpha_H^{-1}(h_{22})g_2\\
=&a_1(\alpha_H^{-2}(h_{1(0)})\cdot\alpha_A^{-1}(b_1))\otimes\alpha_H^{-4}(h_{2111(0)})\alpha_H^{-1}(g_{1(0)})\otimes\{\alpha_A^{-1}(a_2)\alpha_A^{-2}(h_{1(1)})\}\\
&\{[\alpha_A^{-6}(h_{2111(1)})(\alpha_H^{-6}(h_{2112})\cdot\alpha_A^{-3}(b_2))](\alpha_H^{-4}(h_{212})\cdot\alpha_A^{-3}(g_{1(1)}))\}\otimes\alpha_H^{-1}(h_{22})g_2\\
=&a_1(\alpha_H^{-2}(h_{1(0)})\cdot\alpha_A^{-1}(b_1))\otimes\alpha_H^{-4}(h_{2111(0)})\alpha_H^{-1}(g_{1(0)})\otimes\{\alpha_A^{-1}(a_2)\alpha_A^{-2}(h_{1(1)})\}\\
&\{\alpha_A^{-5}(h_{2111(1)})[(\alpha_H^{-6}(h_{2112})\cdot\alpha_A^{-3}(b_2))(\alpha_H^{-5}(h_{212})\cdot\alpha_A^{-4}(g_{1(1)}))]\}\otimes\alpha_H^{-1}(h_{22})g_2\\
=&a_1(\alpha_H^{-2}(h_{1(0)})\cdot\alpha_A^{-1}(b_1))\otimes\alpha_H^{-3}(h_{211(0)})\alpha_H^{-1}(g_{1(0)})\otimes\{\alpha_A^{-1}(a_2)\alpha_A^{-2}(h_{1(1)})\}\\
&\{\alpha_A^{-4}(h_{211(1)})[\alpha_H^{-4}(h_{212})\cdot\alpha_A^{-3}(b_2)\alpha_A^{-4}(g_{1(1)})]\}\otimes\alpha_H^{-1}(h_{22})g_2\\
=&a_1(\alpha_H^{-2}(h_{1(0)})\cdot\alpha_A^{-1}(b_1))\otimes\alpha_H^{-3}(h_{211(0)})\alpha_H^{-1}(g_{1(0)})\\
&\otimes\{\alpha_A^{-1}(a_2)[\alpha_A^{-3}(h_{1(1)})\alpha_A^{-5}(h_{211(1)})]\}\{\alpha_H^{-3}(h_{212})\cdot\alpha_A^{-2}(b_2)\alpha_A^{-3}(g_{1(1)})\}\otimes\alpha_H^{-1}(h_{22})g_2\\
=&a_1(\alpha_H^{-4}(h_{111(0)})\cdot\alpha_A^{-1}(b_1))\otimes\alpha_H^{-3}(h_{112(0)})\alpha_H^{-1}(g_{1(0)})\\
&\otimes\{\alpha_A^{-1}(a_2)\alpha_A^{-5}(h_{111(1)}h_{112(1)})\}\{\alpha_H^{-2}(h_{12})\cdot\alpha_A^{-2}(b_2)\alpha_A^{-3}(g_{1(1)})\}\otimes h_2g_2\\
=&a_1(\alpha_H^{-4}(h_{11(0)1})\cdot\alpha_A^{-1}(b_1))\otimes\alpha_H^{-3}(h_{11(0)2})\alpha_H^{-1}(g_{1(0)})\otimes\{\alpha_A^{-1}(a_2)\alpha_A^{-3}(h_{11(1)})\}\\
&\{\alpha_H^{-2}(h_{12})\cdot\alpha_A^{-2}(b_2)\alpha_A^{-3}(g_{1(1)})\}\otimes h_2g_2\\
=&a_1(\alpha_H^{-3}(h_{1(0)1})\cdot\alpha_A^{-1}(b_1))\otimes\alpha_H^{-2}(h_{1(0)2})\alpha_H^{-1}(g_{1(0)})\otimes\{\alpha_A^{-1}(a_2)\alpha_A^{-2}(h_{1(1)})\}\\
&\{\alpha_H^{-2}(h_{21})\cdot\alpha_A^{-2}(b_2)\alpha_A^{-3}(g_{1(1)})\}\otimes\alpha_H^{-1}(h_{22})g_2\\
=&\Delta(a\otimes h)\Delta(b\otimes g).
\end{aligned}
$$

It is straightforward to check that ε is a morphism of Hom-associate algebra. For the antipode S we have

$$
\begin{aligned}
&(a\otimes h)_1S((a\otimes h)_2)\\
=&(a_1\otimes\alpha_H^{-1}(h_{1(0)}))[(1\otimes S_H(\alpha_H^{-2}(h_{2(0)}))\\
&(S_A((\alpha_A^{-3}(a_2)\alpha_A^{-4}(h_{1(1)}))\alpha_A^{-3}(h_{2(1)}))\otimes 1)]\\
=&[a_1\otimes\alpha_H^{-2}(h_{1(0)})S_H(\alpha_H^{-2}(h_{2(0)}))]\\
&[S_A(\alpha_A^{-1}(a_2)(\alpha_A^{-3}(h_{1(1)})\alpha_A^{-3}(h_{2(1)})))\otimes 1]\\
=&[a_1\otimes\alpha_H^{-2}(h_{(0)1})S_H(\alpha_H^{-2}(h_{(0)2}))][S_A(\alpha_A^{-1}(a_2)(\alpha_A^{-1}(h_{(1)})))\otimes 1]\\
=&[a_1\otimes\varepsilon(h_{(0)})1][S_A(\alpha_A^{-1}(a_2)(\alpha_A^{-1}(h_{(1)})))\otimes 1]\\
=&\varepsilon_A(a)\varepsilon_H(h)1\otimes 1.
\end{aligned}
$$

Similarly we have $S(a\otimes h)_1(a\otimes h)_2=\varepsilon_A(a)\varepsilon_H(h)1\otimes 1$. Thus $A\otimes H$ is a Hom-Hopf algebra.

This finishes the proof. ■

Example 2.7. We now consider A_x^1 given in Example 2.2 (3). By a simple computation the conditions in Theorem 2.6 are satisfied. Hence we have a bicrossproduct Hom-Hopf algebra $(A\#H, \beta\#id)$ with the following Hom-Hopf algebra structure:

The product is given by:

	$1_A\#1_H$	$1_A\#g$	$x\#1_H$	$x\#g$
$1_A\#1_H$	$1_A\#1_H$	$1_A\#g$	$-x\#1_H$	$-x\#g$
$1_A\#g$	$1_A\#g$	$1_A\#1_H$	$x\#g$	$x\#1_H$
$x\#1_H$	$-x\#1_H$	$-x\#g$	0	0
$x\#g$	$-x\#g$	$-x\#1$	0	0

The coproduct is defined as:

$$\Delta(1_A\#1_H) = 1_A\#1_H \otimes 1_A\#1_H,\ \Delta(1_A\#g) = 1_A\#g \otimes 1_A\#g,$$

$$\Delta(x\#1_H) = -x\#1_H \otimes 1_A\#1_H - 1_A\#1_H \otimes x\#1_H,$$

$$\Delta(x \otimes g) = 1_A\#g \otimes x\#g - x\#g \otimes 1_A\#g,$$

and the antipode is given by:

$$S(1_A\#1_H) = 1_A\#1_H,\quad S(1_A\#g) = 1_A\#g,$$

$$S(x\#1_H) = -x\#1_H,\quad S(x\#g) = x\#g.$$

Example 2.8. Let A and H be two Hopf algebras, and $A\#H$ the bicrossproduct of A and H with A a left H-module algebra and H a right A-comodule coalgebra. Assume that α_A and α_H are the automorphisms of Hopf algebras of A and H respectively, and satisfy

$$\alpha_A(h \cdot a) = \alpha_H(h) \cdot \alpha_A(a),$$
$$\alpha_H(h)_{(0)} \otimes \alpha_H(h)_{(1)} = \alpha_H(h_{(0)}) \otimes \alpha_A(h_{(1)}).$$

Then A_{α_A} is a left H_{α_H}-module algebra under the action $h \triangleright a = \alpha_H(h) \cdot \alpha_A(a)$ and H_{α_H} is a right A_{α_A}-comodule coalgebra under the coaction $h_{[0]} \otimes h_{[1]} = \alpha_H(h_{(0)}) \otimes \alpha_A(h_{(1)})$. Hence we have a Hom-type bicrossproduct $A_{\alpha_A}\#H_{\alpha_H}$.

Next we will construct a class of Hom-Hopf algebras.

Corollary 2.9. For any Hom-Hopf algebra (H, S, α), there exists a bicrossproduct structure on the space $(H \otimes H^{op}, \alpha \otimes \alpha)$.

Proof. From the Example 2.5 and Proposition 2.4, (H^{op}, α) is a right (H, α)-comodule Hom-coalgebra. And (H, α) is a left (H^{op}, α)-module Hom-algebra under the action

$$h \cdot a = (S(\alpha^{-2}(h_1))\alpha^{-1}(a))\alpha^{-1}(h_2),$$

for any $a, h \in H$. The verification of module condition is straightforward and is left to the reader.

For any $h, a, b \in H$,

$$\begin{aligned}
&(h_1 \cdot a)(h_2 \cdot b)\\
=&[(S(\alpha^{-2}(h_{11}))\alpha^{-1}(a))\alpha^{-1}(h_{12})][(S(\alpha^{-2}(h_{21}))\alpha^{-1}(b))\alpha^{-1}(h_{22})]\\
=&[(S(\alpha^{-2}(h_{11}))\alpha^{-1}(a))(\alpha^{-2}(h_{12})S(\alpha^{-2}(h_{21}))][b\alpha^{-1}(h_{22})]\\
=&[(S(\alpha^{-1}(h_1))\alpha^{-1}(a))(\alpha^{-3}(h_{211})S(\alpha^{-2}(h_{212}))][b\alpha^{-1}(h_{22})]\\
=&(S(h_1)a)(bh_2) = (S(h_1)\alpha^{-1}(ab))\alpha(h_2)\\
=&\alpha^2(h) \cdot (ab).
\end{aligned}$$

That is, (H, α) is a left (H^{op}, α)-module Hom-algebra.

Moreover for any $h, g, a \in H$,

$$\begin{aligned}
&(h \cdot a)_1 \otimes (h \cdot a)_2\\
=&(S(\alpha^{-2}(h_{12}))\alpha^{-1}(a_1))\alpha^{-1}(h_{21}) \otimes (S(\alpha^{-2}(h_{11}))\alpha^{-1}(a_2))\alpha^{-1}(h_{22})\\
=&(S(\alpha^{-3}(h_{121}))\alpha^{-1}(a_1))\alpha^{-2}(h_{122}) \otimes [S(\alpha^{-2}(h_{11}))(\alpha^{-4}(h_{211})S\alpha^{-4}(h_{212}))]\\
&(\alpha^{-1}(a_2)\alpha^{-2}(h_{22}))\\
=&(S(\alpha^{-4}(h_{1121}))\alpha^{-1}(a_1))\alpha^{-3}(h_{1122}) \otimes [S(\alpha^{-3}(h_{111}))\alpha^{-2}(h_{12})]\\
&[(S\alpha^{-3}(h_{21})\alpha^{-2}(a_2))\alpha^{-2}(h_{22})]\\
=&(S(\alpha^{-3}(h_{1(0)1}))\alpha^{-1}(a_1))\alpha^{-2}(h_{1(0)2}) \otimes \alpha^{-1}(h_{1(1)})[(S\alpha^{-3}(h_{21})\alpha^{-2}(a_2))\alpha^{-2}(h_{22})]\\
=&\alpha^{-1}(h_{1(0)} \cdot a_1 \otimes \alpha^{-1}(h_{1(1)})(\alpha^{-1}(h_2) \cdot \alpha^{-1}(b_2)),
\end{aligned}$$

and

$$\begin{aligned}
&(gh)_0 \otimes (gh)_1 = \alpha^{-1}(g_{12}h_{12}) \otimes S(\alpha^{-2}(g_{11}h_{11}))\alpha^{-1}(g_2h_2)\\
=&\alpha^{-1}(g_{12})\alpha^{-1}(h_{12}) \otimes [S(\alpha^{-2}(h_{11}))S(\alpha^{-2}(g_{11}))][\alpha^{-1}(g_2)\alpha^{-1}(h_2)]\\
=&\alpha^{-1}(g_{12})\alpha^{-1}(h_{12}) \otimes S(\alpha^{-1}(h_{11}))[(S(\alpha^{-3}(g_{11}))\alpha^{-2}(g_2))\alpha^{-1}(h_2)]\\
=&\alpha^{-1}(g_{12})\alpha^{-2}(h_{112}) \otimes [S(\alpha^{-3}(h_{111}))\alpha^{-2}(h_{12})]\\
&[(S(\alpha^{-3}(h_{21}))(S(\alpha^{-4}(g_{11}))\alpha^{-3}(g_2)))\alpha^{-2}(h_{22})]\\
=&g_{(0)}\alpha^{-1}(h_{1(0)}) \otimes \alpha^{-1}(h_{1(1)})(\alpha^{-1}(h_2) \cdot \alpha^{-1}(g_{(1)})),
\end{aligned}$$

Finally

$$\begin{aligned}
&h_{2(0)} \otimes (h_1 \cdot a)h_{2(1)} \\
=&\alpha^{-1}(h_{212}) \otimes [(S(\alpha^{-2}(h_{11}))\alpha^{-1}(a))\alpha^{-1}(h_{12})][S(\alpha^{-2}(h_{211}))\alpha^{-1}(h_{22})] \\
=&\alpha^{-1}(h_{212}) \otimes [(S(\alpha^{-2}(h_{11}))\alpha^{-1}(a))(\alpha^{-2}(h_{12})S(\alpha^{-3}(h_{211})))]h_{22} \\
=&\alpha^{-1}(h_{122}) \otimes [(S(\alpha^{-2}(h_{11}))\alpha^{-1}(a))(\alpha^{-4}(h_{1211})S(\alpha^{-4}(h_{1212})))]\alpha(h_2) \\
=&h_{12} \otimes (S(\alpha^{-1}(h_{11}))a)\alpha(h_2) \\
=&h_{12} \otimes S(h_{11})(ah_2) \\
=&h_{12} \otimes [S(\alpha^{-1}(h_{11}))(\alpha^{-3}(h_{211})S(\alpha^{-3}(h_{212}))](a\alpha^{-1}(h_{22})) \\
=&h_{1(0)} \otimes h_{1(1)}(h_2 \cdot a).
\end{aligned}$$

This finishes the proof. ■

Therefore by Theorem 2.6, we have a bicrossproduct structure on $H \otimes H^{op}$ with multiplication and comultiplication as follows:

$$(a \ltimes h)(b \ltimes k) = a[(S(\alpha^{-4}(h_{11}))\alpha^{-2}(b))\alpha^{-3}(h_{12})] \ltimes k\alpha^{-1}(h_2),$$

$$\Delta(a \ltimes h) = a_1 \ltimes \alpha^{-2}(h_{112}) \otimes \alpha^{-1}(a_2)(S(\alpha^{-4}(h_{111}))\alpha^{-3}(h_{12})) \ltimes h_2,$$

for any $a, b, h, k \in H$.

Definition 2.10. Let (H, α_H) be a Hom-bialgebra and (C, α_C) a Hom-coassociative coalgebra. Then C is called a left *H-module Hom-coalgebra* if C is a left H-module with the action $H \otimes C \longrightarrow C,\ h \otimes c \mapsto h \cdot c$, such that

$$\Delta_C(h \cdot c) = h_1 \cdot c_1 \otimes h_2 \cdot c_2, \quad \varepsilon_C(h \cdot c) = \varepsilon_H(h)\varepsilon_C(c),$$

for any $h \in H,\ c \in C$.

Definition 2.11. Let (A, α_A) and (H, α_H) be two Hom-bialgebras. (A, H) is called a *matched pair* if there exist linear maps

$$\triangleleft : H \otimes A \longrightarrow H,\ \triangleright : H \otimes A \longrightarrow A,$$

turning A into a left H-module Hom-coalgebra and turning H into a right A-module Hom-coalgebra such that the following conditions are satisfied:

$$(hg) \triangleleft a = (h \triangleleft (\alpha_H^{-2}(g_1) \triangleright \alpha_A^{-3}(a_1))(\alpha_H^{-1}(g_2) \triangleleft \alpha_A^{-2}(a_2)) \tag{2. 5}$$

$$h \triangleright (ab) = (\alpha_H^{-2}(h_1) \triangleright \alpha_A^{-1}(a_1))((\alpha_H^{-3}(h_2) \triangleleft \alpha_A^{-2}(a_2)) \triangleright b), \tag{2. 6}$$

$$h_1 \triangleleft a_1 \otimes h_2 \triangleright a_2 = h_2 \triangleleft a_2 \otimes h_1 \triangleright a_1. \tag{2. 7}$$

Proposition 2.12. Let (A, S_A, α_A) and (H, S_H, α_H) be two Hom-Hopf algebras, and (A, H) a matched pair. There exists a unique Hom-Hopf algebra structure on the vector space $A \otimes H$ with the multiplication, comultiplication and the antipode given by

$$\begin{aligned}
&(a \otimes h)(b \otimes g) = a(\alpha_H^{-2}(h_1) \triangleright \alpha_A^{-2}(b_1)) \otimes (\alpha_H^{-2}(h_2) \triangleleft \alpha_A^{-2}(b_2))g,\\
&\Delta(a \otimes h) = a_1 \otimes h_1 \otimes a_2 \otimes h_2,\\
&S(a \otimes h) = (1_A \otimes S_H\alpha_H^{-1}(h))(S_A\alpha_A^{-1}(a) \otimes 1_H),
\end{aligned}$$

for any $a, b \in A$ and $g, h \in H$.

Equipped with this Hopf algebra structure, $A \otimes H$ is called double crossed product of A and H denoted by $A \bowtie H$.

Proof. Define the linear map $T : H \otimes A \longrightarrow A \otimes H$ by

$$T(h \otimes a) = \alpha_H^{-2}(h_1) \triangleright \alpha_A^{-2}(b_1) \otimes \alpha_H^{-2}(h_2) \triangleleft \alpha_A^{-2}(b_2),$$

for any $a \in A$ and $h \in H$.

By Proposition 2.6 in [8], in order to prove the multiplication is Hom-associative, we need only to verify that T is a Hom-twisting map between H and A. Indeed, first easy to see that

$$T \circ (\alpha_H \otimes \alpha_A) = (\alpha_A \otimes \alpha_H) \circ T.$$

Then for any $a, b \in A$ and $h \in H$,

$$\begin{aligned}
&(ab)_T \otimes \alpha_H(h)_T\\
=&\alpha_H^{-1}(h_1) \triangleright \alpha_A^{-2}(a_1b_1) \otimes \alpha_H^{-1}(h_2) \triangleleft \alpha_A^{-2}(a_2b_2)\\
=&\alpha_H^{-1}(h_1) \triangleright \alpha_A^{-2}(a_1b_1) \otimes \alpha_H^{-1}(h_2) \triangleleft \alpha_A^{-2}(a_2b_2)\\
=&(\alpha_H^{-3}(h_{11}) \triangleright \alpha_A^{-3}(a_{11}))((\alpha_H^{-4}(h_{12}) \triangleleft \alpha_A^{-4}(a_{12})) \triangleright \alpha^{-2}(b_1)) \otimes \alpha_H^{-1}(h_2) \triangleleft \alpha_A^{-2}(a_2b_2)\\
=&(\alpha_H^{-2}(h_1) \triangleright \alpha_A^{-2}(a_1))((\alpha_H^{-4}(h_{21}) \triangleleft \alpha_A^{-4}(a_{21})) \triangleright \alpha^{-2}(b_1))\\
&\otimes (\alpha_H^{-3}(h_{22}) \triangleleft \alpha_A^{-3}(a_{22})) \triangleleft \alpha_A^{-1}(b_2)\\
=&a_Tb_t \otimes \alpha_H(h_{Tt}),
\end{aligned}$$

where we have used the notation $T(h \otimes a) = a_T \otimes h_T = a_t \otimes h_t$.

Similarly for any $a \in A$ and $g, h \in H$, we have

$$\alpha_A(a)_T \otimes (hg)_T = \alpha_A(a_{Tt}) \otimes h_tg_T.$$

Now $A \otimes H$ is a Hom-associative algebra. Next for any $a, b \in A, h, g \in H$,

$$
\begin{aligned}
&\Delta((a\otimes h)(b\otimes g))\\
=&\Delta(a(\alpha_H^{-2}(h_1)\triangleright\alpha_A^{-2}(b_1))\otimes\alpha_H^{-2}(h_2)\triangleleft\alpha_A^{-2}(b_2))g\\
=&a_1(\alpha_H^{-2}(h_{11})\triangleright\alpha_A^{-2}(b_{11}))\otimes(\alpha_H^{-2}(h_{21})\triangleleft\alpha_A^{-2}(b_{21}))g_1\\
&\otimes a_2(\alpha_H^{-2}(h_{12})\triangleright\alpha_A^{-2}(b_{12}))\otimes(\alpha_H^{-2}(h_{22})\triangleleft\alpha_A^{-2}(b_{22}))g_2\\
=&a_1(\alpha_H^{-2}(h_{11})\triangleright\alpha_A^{-2}(b_{11}))\otimes(\alpha_H^{-2}(h_{12})\triangleleft\alpha_A^{-2}(b_{12}))g_1\\
&\otimes a_2(\alpha_H^{-2}(h_{21})\triangleright\alpha_A^{-2}(b_{21}))\otimes(\alpha_H^{-2}(h_{22})\triangleleft\alpha_A^{-2}(b_{22}))g_2\\
=&\Delta(a\otimes h)\Delta(b\otimes g).
\end{aligned}
$$

Therefore $A\otimes H$ is a Hom-bialgebra. It is straightforward to check that S is the antipode.

This finishes the proof. ■

Let (H,α) be a Hom-bialgebra. In [8], the authors have defined the Hom-associative algebra $(H^*,(\alpha^{-1})^*)$, the linear dual of H, where the multiplication is given by

$$(f\bullet g)(h)=f(\alpha^{-2}(h_1))g(\alpha^{-2}(h_2)),$$

for any $f,g\in H^*$, and $h\in H$.

Now we give the comultiplication on H^* by

$$f(gh)=f_1(\alpha^2(h))f_2(\alpha^2(g)),$$

for any $f\in H^*$ and $g,h\in H$. In other words, $\langle\Delta(f),h\otimes k\rangle=f(\alpha^{-2}(hk))$.

Proposition 2.13. Let (H,S,α) be a Hom-Hopf algebra. With the multiplication and comultiplication defined on H^* as above, $(H^*,S^*,(\alpha^{-1})^*)$ is a Hom-Hopf algebra.

Proof First we need to prove that $(H^*,(\alpha^{-1})^*)$ is a Hom-coassociative coalgebra. Indeed for any $g,h,k\in H$ and $f\in H^*$,

$$
\begin{aligned}
f((gh)\alpha(k))&=f_1(\alpha^2(gh))f_2(\alpha^3(k))\\
&=f_{11}(\alpha^4(g))f_{12}(\alpha^4(h))f_2(\alpha^3(k)),
\end{aligned}
$$

and

$$
\begin{aligned}
f(\alpha(g)(hk))&=f_1(\alpha^3(g))f_2(\alpha^2(hk))\\
&=f_1(\alpha^3(g))f_{21}(\alpha^4(h))f_{22}(\alpha^4(k)),
\end{aligned}
$$

therefore $(\alpha^{-1})^*(f_1)\otimes f_{21}\otimes f_{22}=f_{11}\otimes f_{12}\otimes(\alpha^{-1})^*(f_2)$.

Furthermore

$$\langle\Delta\circ(\alpha^{-1})^*(f),h\otimes k\rangle=(\alpha^{-1})^*(f)(\alpha^{-2}(hk))=f(\alpha^{-3}(hk)),$$

and

$$(\alpha^{-1})^*(f_1)(h)(\alpha^{-1})^*(f_2)(k)=f_1(\alpha^{-1}(h))f_2(\alpha^{-1}(k))=f(\alpha^{-3}(hk)).$$

That is, $\Delta \circ (\alpha^{-1})^* = ((\alpha^{-1})^* \otimes (\alpha^{-1})^*) \circ \Delta$. Then $(H^*, (\alpha^{-1})^*)$ is a Hom-coassociative coalgebra.

Finally for any $f, g \in H^*$ and $h, k \in H$,

$$\begin{aligned}\langle \Delta(f \bullet g), h \otimes k\rangle &= (f \bullet g)(\alpha^{-2}(hk)) \\ &= f(\alpha^{-4}(h_1k_1))g(\alpha^{-4}(h_2k_2)) \\ &= f_1(\alpha^{-2}(h_1))f_2(\alpha^{-2}(k_1))g_1(\alpha^{-2}(h_2))g_2(\alpha^{-2}(k_2)) \\ &= \langle f_1 \bullet g_1 \otimes f_2 \bullet g_2, h \otimes k\rangle.\end{aligned}$$

Therefore $(H^*, (\alpha^{-1})^*)$ is a Hom-bialgebra. It is not hard to verify that S^* is the antipode.

This finishes the proof. ■

The following proposition can be found in [8].

Proposition 2.14. Let (C, α_C) be a Hom-coassociative coalgebra, and (M, α_M) is a right C-comodule. Then (M, α_M) is a left $(C^*, (\alpha_C^{-1})^*)$-module with the action $f \cdot m = f(m_{(1)})m_{(0)}$.

Proposition 2.15. Let (A, α_A) and (H, α_H) be two Hom-bialgebras, $(A \ltimes H, \alpha_A \otimes \alpha_H)$ the bicrossproduct of A and H. Define the left action $\triangleright : A^* \otimes H \longrightarrow H$ of A^* on H by

$$f \triangleright h = f(h_{(1)})h_{(0)},$$

and the right action $\triangleleft : A^* \otimes H \longrightarrow A^*$ of H on A^* by

$$\langle f \triangleleft h, a\rangle = \langle f, h \cdot \alpha^{-2}(a)\rangle,$$

for any $f \in A^*$, $h \in H$ and $a \in A$. Then $(A^*, (\alpha_A^{-1})^*)$ and (H, α_H) is a matched pair.

Proof. First, we need to verify that H is a left A^*-module Hom-coalgebra. By Proposition 2.14, H is a left A^*-module. Then for any $h \in H$ and $f \in A^*$,

$$\begin{aligned}\Delta(f \triangleright h) &= f(h_{(1)})h_{(0)1} \otimes h_{(0)2} \\ &= f(\alpha^{-2}(h_{1(1)}h_{2(1)}))h_{1(0)} \otimes h_{2(0)} \\ &= f_1(h_{1(1)})h_{1(0)} \otimes f_2(h_{2(1)})h_{2(0)} \\ &= f_1 \triangleright h_1 \otimes f_2 \triangleright h_2.\end{aligned}$$

So H is a left A^*-module Hom-coalgebra.

Now for any $g, h \in H$, $f \in A^*$ and $a, b \in A$

$$\begin{aligned}\langle (\alpha_A^{-1})^*(f) \triangleleft (hg), a\rangle &= \langle f, \alpha_H^{-1}(hg) \cdot \alpha_A^{-3}(a)\rangle \\ &= \langle f \triangleleft h, \alpha_H(g) \cdot \alpha_A^{-2}(a)\rangle \\ &= \langle (f \triangleleft h) \triangleleft \alpha_H(g), a\rangle.\end{aligned}$$

and obviously

$$(\alpha_A^{-1})^*(f \triangleleft h) = (\alpha_A^{-1})^*(f) \triangleleft \alpha_H(h).$$

Then

$$\begin{aligned}\langle \Delta(f \triangleleft h), a \otimes b\rangle &= \langle f \triangleleft h, \alpha_A^{-2}(ab)\rangle = \langle f, h \cdot \alpha_A^{-4}(ab)\rangle \\ &= \langle f, (\alpha_H^{-2}(h_1) \cdot \alpha_A^{-4}(a))(\alpha_H^{-2}(h_2) \cdot \alpha_A^{-4}(b))\rangle \\ &= \langle f_1, h_1 \cdot \alpha_A^{-2}(a)\rangle\langle f_2, h_2 \cdot \alpha_A^{-2}(b)\rangle \\ &= \langle f_1 \triangleleft h_1, a\rangle\langle f_2 \triangleleft h_2, b\rangle.\end{aligned}$$

Therefore A^* is a right H-module Hom-coalgebra.

Next we will verify the compatible conditions in Definition 2.11. For any $f, g \in A^*$, $a \in A$ and $h, k \in H$,

$$\begin{aligned}&\langle (f \bullet g) \triangleleft h,\ a\rangle = \langle f \bullet g,\ h \cdot \alpha_A^{-2}(a)\rangle \\ =&\langle f,\ \alpha_A^{-2}((h \cdot \alpha_A^{-2}(a))_1)\rangle\langle g,\ \alpha_A^{-2}((h \cdot \alpha_A^{-2}(a))_2)\rangle \\ =&\langle f,\ \alpha_H^{-3}(h_{1(0)}) \cdot \alpha_A^{-4}(a_1)\rangle\langle g,\ \alpha_H^{-3}(h_{1(1)})(\alpha_H^{-3}(h_2) \cdot \alpha_A^{-5}(a_2))\rangle \\ =&\langle f,\ \alpha_H^{-3}(h_{1(0)}) \cdot \alpha_A^{-4}(a_1)\rangle\langle g_1,\ \alpha_H^{-1}(h_{1(1)})\rangle\langle g_2,\ \alpha_H^{-1}(h_2) \cdot \alpha_A^{-3}(a_2)\rangle \\ =&\langle f,\ ((\alpha_A^*)^2(g_1) \triangleright \alpha_H^{-3}(h_1)) \cdot \alpha_A^{-4}(a_1)\rangle\langle \alpha_A^*(g_2),\ \alpha_H^{-2}(h_2) \cdot \alpha_A^{-4}(a_2)\rangle \\ =&\langle f \triangleleft ((\alpha_A^*)^2(g_1) \triangleright \alpha_H^{-3}(h_1)),\ \alpha_A^{-2}(a_1)\rangle\langle \alpha_A^*(g_2) \triangleleft \alpha_H^{-2}(h_2),\ \alpha_A^{-2}(a_2)\rangle \\ =&\langle [f \triangleleft ((\alpha_A^*)^2(g_1) \triangleright \alpha_H^{-3}(h_1))] \bullet [\alpha_A^*(g_2) \triangleleft \alpha_H^{-2}(h_2)], a\rangle,\end{aligned}$$

thus we have the condition (2.5). And by

$$\begin{aligned}f \triangleright (hk) &= f((hk)_{(1)})(hk)_{(0)} \\ &= \langle f,\ \alpha_A^{-1}(h_{1(1)})(\alpha_H^{-1}(h_2) \cdot \alpha_A^{-1}(k_{(1)}))\rangle\alpha_H^{-1}(h_{1(0)})k_{(0)} \\ &= \langle f_1,\ \alpha_A(h_{1(1)})\rangle\langle f_2,\ \alpha_H(h_2) \cdot \alpha_A(k_{(1)})\rangle\alpha_H^{-1}(h_{1(0)})k_{(0)} \\ &= \langle (\alpha_A^*)^2(f_1),\ \alpha_A^{-1}(h_{1(1)})\rangle\langle (\alpha_A^*)^3(f_2) \triangleleft \alpha_H^{-2}(h_2),\ k_{(1)}\rangle\alpha_H^{-1}(h_{1(0)})k_{(0)} \\ &= [(\alpha_A^*)^2(f_1) \triangleright \alpha_H^{-1}(h_1)][((\alpha_A^*)^3(f_2) \triangleleft \alpha_H^{-2}(h_2)) \triangleright k],\end{aligned}$$

we get the condition (2.6). As for the condition (2.7),

$$\begin{aligned}\langle f_1 \triangleleft h_1,\ a\rangle\langle g,\ f_2 \triangleright h_2\rangle &= \langle f_1,\ h_1 \cdot \alpha_A^{-2}(a)\rangle\langle g,\ h_{2(0)}\rangle\langle f_2,\ h_{2(1)}\rangle \\ &= \langle f,\ \alpha_A^{-2}((h_1 \cdot \alpha_A^{-2}(a))h_{2(1)})\rangle\langle g,\ h_{2(0)}\rangle \\ &= \langle f,\ \alpha_A^{-2}(h_{1(1)}(h_2 \cdot \alpha_A^{-2}(a)))\rangle\langle g,\ h_{1(0)}\rangle \\ &= \langle f_1,\ h_{1(1)}\rangle\langle f_2,\ h_2 \cdot \alpha_A^{-2}(a)\rangle\langle g,\ h_{1(0)}\rangle \\ &= \langle f_2 \triangleleft h_2,\ a\rangle\langle g,\ f_1 \triangleright h_1\rangle.\end{aligned}$$

Therefore $(A^*, (\alpha_A^{-1})^*)$ and (H, α_H) is a matched pair.

This finishes the proof. ■

By this result, we have the double crossed product $(H \bowtie A^*, \alpha_H \otimes (\alpha_A^{-1})^*)$ with the multiplication and comultiplication given by

$$(h \otimes f)(k \otimes g) = h\alpha_H^{-2}(k_{1(0)}) \otimes \langle f,\ \alpha_A^{-2}(k_{1(1)})(\alpha_H^{-2}(k_2) \cdot \alpha_A^{-2}(?))\rangle g,$$

$$\Delta(h \otimes f) = h_1 \otimes f_1 \otimes h_2 \otimes f_2,$$

for any $h, k \in H$ and $f, g \in A^*$.

Corollary 2.16. Let (H, α) be a Hom-Hopf algebra. We have the Drinfeld double $(H^{op} \bowtie H^*, \alpha \otimes (\alpha^{-1})^*)$ with the tensor comultiplication and multiplication given by

$$(h \otimes f)(k \otimes g) = \alpha^{-2}(k_{21})h \otimes [\alpha^{-3}(k_{22}) \rightharpoonup ((\alpha^*)^2(f) \leftharpoonup S\alpha^{-3}(k_1))]g,$$

for any $h, k \in H$ and $f, g \in H^*$, where $\langle f \leftharpoonup h,\ k\rangle = \langle f,\ h\alpha^{-2}(k)\rangle$ and $\langle h \rightharpoonup f,\ k\rangle = \langle f,\ \alpha^{-2}(k)h\rangle$.

Example 2.17. (1) Let G be a finite group and ϕ an automorphism of G. Then (kG, ϕ) with the following structure is a Hom-Hopf algebra:

$$g \cdot h = \phi(gh),\ \Delta(g) = \phi(g) \otimes \phi(g),\ \varepsilon(g) = 1,\ S(g) = g^{-1}.$$

Let $\{e_g\}_{g \in G}$ be the dual basis of the basis of kG. Then we have the Hom-Hopf algebra k^G, dual of kG, with the multiplication

$$e_g e_h = \delta_{g,h}\ e_{\phi(g)},$$

and the comultiplication, counit and antipode

$$\Delta^*(e_g) = \sum_{uv=\phi(g)} e_u \otimes e_v,\ \varepsilon^*(e_g) = \delta_{g,1},\ S^*(e_g) = e_{g^{-1}},$$

for any $g, h \in G$.

By Corollary 2.16, the multiplication in $D(kG)$ is given by

$$(g \otimes e_h)(p \otimes e_q) = \sum \phi(pg) \otimes \delta_{\phi(p)h\phi(p^{-1}),q}\ e_{\phi(q)}.$$

(2) In particular let $G = (g)$ be a cycle group of order $n > 0$. Define $\phi \in Aut(G)$ by

$$\phi(g^i) = g^{-i}, \quad 0 \le i < n.$$

Obviously $\phi \in Aut(G)$. Then we have a Hom-Hopf algebra (kG, ϕ) by

$$g^i \cdot g^j = g^{n-(i+j)},\ \Delta(g^i) = g^{n-i} \otimes g^{n-i},\ \varepsilon(g^i) = 1,\ S(g^i) = g^{n-i},$$

for any $0 \leq i, j < n$. Let $\{e_i\}_{0 \leq i < n}$ be the dual basis of kG such that $e^i(g^j) = \delta_{i,j} e_i$. Then

$$\Delta^*(e_i) = \sum_{j+k=n-i} e_j \otimes e_k,\ \varepsilon^*(e_i) = \delta_{i,1},\ S^*(e_i) = e_{n-i}.$$

In the Drinfeld double $D(kG)$, we have the product

$$(g^i \otimes e_j)(g^m \otimes e_k) = \sum g^{n-(i+m)} \otimes \delta_{j,k}\ e_{n-k},$$

for any $0 \leq i, j, m, k < n$.

Example 2.18. Let H be the Hom-algebra generated by the elements $1_H,\ g$ and x satisfying the following relations:

$$1_H 1_H = 1_H,\ 1_H g = g 1_H = g,\ 1_H x = x 1_H = -x,$$

$$g^2 = 1_H,\ x^2 = 0,\ gx = -xg.$$

The automorphism $\alpha : H \longrightarrow H$ is defined by

$$\alpha(1_H) = 1_H,\ \alpha(g) = g,\ \alpha(x) = -x,\ \alpha(gx) = -gx.$$

Then (H, α) is a Hom-associative algebra, and $\alpha^2 = id$.

Define

$$\Delta(1_H) = 1_H \otimes 1_H,\ \Delta(g) = g \otimes g,$$

$$\Delta(x) = (-x) \otimes g + 1 \otimes (-x),$$

$$\varepsilon(1_H) = 1,\ \varepsilon(g) = 1,\ \varepsilon(x) = 0,$$

$$S(1_H) = 1_H,\ S(g) = g,\ S(x) = -gx.$$

Then (H, α) is a monoidal Hom-Hopf algebra.

Furthermore, let $R = \frac{1}{2}(1 \otimes 1 + 1 \otimes g + g \otimes 1 - g \otimes g)$. Then it is not hard to check that (H, α, R) is quasitriangular.

Proposition 2.19. The Drinfeld double $(H^{op} \bowtie H^*, \alpha \otimes (\alpha^{-1})^*)$ has the quasitriangular structure

$$R = 1 \bowtie (\alpha^{-1})^*(e^i) \otimes S^{-1}(e_i) \bowtie \varepsilon,$$

where $\{e_i\}$ and $\{e_i^*\}$ are a base of H and its dual base in H^* respectively.

Proof. For any $h, k, k' \in H$ and $f, g, g' \in H^*$, on one hand,

$$\begin{aligned}
&\langle \Delta^{op}(h \otimes f)R, g \otimes k \otimes g' \otimes k' \rangle \\
=&\langle (h_2 \otimes f_2)(1 \otimes (\alpha^{-1})^*(e^i)), g \otimes k \rangle \langle (h_1 \otimes f_1)(S^{-1}(e_i) \otimes \varepsilon), g' \otimes k' \rangle \\
=&\langle g, \alpha(h_2) \rangle \langle f_2, \alpha^{-2}(k_1) \rangle \langle g', \ S^{-1}(\alpha^{-5}(k_{221}))h_1 \rangle \\
&\langle f_1, \alpha^{-5}(k_{222}))(\alpha^{-3}(k')S^{-1}(\alpha^{-5}(k_{21}))) \rangle \\
=&\langle g, \alpha(h_2) \rangle \langle g', \ S^{-1}(\alpha^{-4}(k_{21}))h_1 \rangle \\
&\langle f_1, \alpha^{-4}(k_{22}))(\alpha^{-3}(k')S^{-1}(\alpha^{-5}(k_{12}))) \rangle \langle f_2, \alpha^{-3}(k_{11}) \rangle \\
=&\langle g, \alpha(h_2) \rangle \langle g', \ S^{-1}(\alpha^{-4}(k_{21}))h_1 \rangle \\
&\langle f, [\alpha^{-6}(k_{22}))(\alpha^{-5}(k')S^{-1}(\alpha^{-7}(k_{12}))]\alpha^{-5}(k_{11}) \rangle \\
=&\langle g, \alpha(h_2) \rangle \langle g', \ S^{-1}(\alpha^{-4}(k_{21}))h_1 \rangle \\
&\langle f, [\alpha^{-6}(k_{22})\alpha^{-4}(k')][S^{-1}(\alpha^{-6}(k_{12}))\alpha^{-6}(k_{11})] \rangle \\
=&\langle g, \alpha(h_2) \rangle \langle g', \ S^{-1}(\alpha^{-3}(k_1))h_1 \rangle \langle f, [\alpha^{-4}(k_2)\alpha^{-3}(k')] \rangle.
\end{aligned}$$

On the other hand,

$$\begin{aligned}
&\langle R\Delta(h \otimes f), g \otimes k \otimes g' \otimes k' \rangle \\
=&\langle (1 \otimes (\alpha^{-1})^*(e^i))(h_1 \otimes f_1), g \otimes k \rangle \langle (S^{-1}(\alpha(e_i)) \otimes \varepsilon)(h_2 \otimes f_2), g' \otimes k' \rangle \\
=&\langle g, \alpha^{-1}(h_{112}) \rangle \langle e^i, S\alpha^{-3}(h_{111})(\alpha^{-5}(k_1)\alpha^{-3}(h_{12})) \rangle \langle f_1, \alpha^{-2}(k_2) \rangle \\
&\langle g', h_2 S^{-1}(\alpha(e_i)) \rangle \langle f_2, \alpha^{-1}(k') \rangle \\
=&\langle g, \alpha^{-1}(h_{112}) \rangle \langle f, \alpha^{-4}(k_2)\alpha^{-3}(k') \rangle \langle g', h_2[S^{-1}(\alpha^{-5}(k_1)\alpha^{-3}(h_{12}))\alpha^{-3}(h_{111})] \rangle \\
=&\langle g, h_{12} \rangle \langle f, \alpha^{-4}(k_2)\alpha^{-3}(k') \rangle \\
&\langle g', [\alpha^{-2}(h_{22})S^{-1}(\alpha^{-2}(h_{21}))][S^{-1}(\alpha^{-4}(k_1))\alpha^{-2}(h_{11})] \rangle \\
=&\langle g, \alpha(h_2) \rangle \langle f, \alpha^{-4}(k_2)\alpha^{-3}(k') \rangle \langle g', [S^{-1}(\alpha^{-3}(k_1))h_1] \rangle.
\end{aligned}$$

Hence $\Delta^{op}(h \otimes f)R = R\Delta(h \otimes f)$. Similarly we have $(\Delta \otimes \alpha)R = R^{13}R^{23}$, $(\alpha \otimes \Delta)R = R^{13}R^{12}$.

This finishes the proof. ■

Example 2.20. In the Example 2.17, the Drinfeld double of (kG, ϕ) is given where G is a finite group and ϕ is a group automorphism of G. Then by the above proposition, the quasitriangular structure of $D(kG)$ is

$$R = \sum_{g \in G} 1_G \bowtie e_{\phi(g)} \otimes g^{-1} \bowtie 1_{k^G}.$$

Especially when G is a cycle group of order n, the quasitriangular structure of $D(kG)$ is

$$R = \sum_{0 \le i < n} 1_G \bowtie e_{n-i} \otimes g^{-i} \bowtie 1_{k^G}.$$

3. DUAL PAIRS OF HOM-HOPF-ALGEBRAS

In this section, we will consider dual pairs of Hom-Hopf algebras and then give the Drinfel'd double associated to a pairing of Hom-Hopf algebras in the setting of dual pairs.

Definition 3.1. Let (A, α_A) and (B, α_B) be two Hom-Hopf algebras and $(-,-) : A \otimes B \longrightarrow k$ a non-degenerate bilinear form. Then (A, B) is called a *dual pair* if the following conditions hold:

$$(a, 1) = \varepsilon(a), \qquad (1, b) = \varepsilon(b), \tag{3. 1}$$

$$(\alpha_A(a), \alpha_B(b)) = (a, b), \tag{3. 2}$$

$$(aa', b) = (\alpha_A^2(a), b_1)(\alpha_A^2(a'), b_2), \tag{3. 3}$$

$$(a, bb') = (a_1, \alpha_B^2(b))(a_2, \alpha_B^2(b')), \tag{3. 4}$$

$$(S_A(a), b) = (a, S_B^{-1}(b)), \tag{3. 5}$$

for any $a, a' \in A$ and $b, b' \in B$.

Example 3.2. Let (A, B, θ) be a dual pair of Hopf algebras defined in [11], α_A and α_B be a Hopf algebra isomorphism of A and B respectively. Assume that $(\alpha_A, \alpha_B, \theta)$ is a compatible pairing in the sense that

$$\theta(\alpha_A(a), \alpha_B(b)) = \theta(a, b),$$

for any $a \in A$ and $b \in B$. Define a bilinear form on $A_{\alpha_A} \otimes B_{\alpha_B}$ by

$$(\alpha_A(a), \alpha_B(b)) = \theta(a, b),$$

then A_{α_A} and B_{α_B} is a dual pair of Hom-Hopf algebra.

Recall from [7] that a Hom-twisting map between the Hom-associative algebras (A, μ_A, α_A) and (B, μ_B, α_B) is a linear map $T : B \otimes A \longrightarrow A \otimes B$ such that the following conditions are satisfied:

$$(\alpha_A \otimes \alpha_B) \circ R = R \circ (\alpha_B \otimes \alpha_A),$$
$$T \circ (\mu_B \otimes \alpha_A) = (\alpha_A \otimes \mu_B) \circ (T \otimes id) \circ (id \otimes T),$$
$$T \circ (\alpha_B \otimes \mu_A) = (\mu_A \otimes \alpha_B) \circ (id \otimes T) \circ (T \otimes id).$$

Then we will construct a Hom-twisting map. In fact, we have two linear maps $R_1, R_2 : A \otimes B \longrightarrow A \otimes B$ given by

$$R_1(a \otimes b) = (\alpha_A(a_2), b_1)\alpha_A^{-1}(a_1) \otimes \alpha_B^{-1}(b_2),$$

$$R_2(a \otimes b) = (\alpha_A(a_1), b_2)\alpha_A^{-1}(a_2) \otimes \alpha_B^{-1}(b_1).$$

Easy to check that both are invertible with the inverses

$$R_1^{-1}(a \otimes b) = (S^{-1}\alpha_A(a_2), b_1)\alpha_A^{-1}(a_1) \otimes \alpha_B^{-1}(b_2),$$

$$R_2^{-1}(a \otimes b) = (S^{-1}\alpha_A(a_1), b_2)\alpha_A^{-1}(a_2) \otimes \alpha_B^{-1}(b_1).$$

Define the linear map $T = R_1 \circ R_2^{-1} \circ \tau : B \otimes A \longrightarrow A \otimes B$, where τ is the flip map.

Proposition 3.3. The above T is a Hom-twisting map.

Proof. Obviously $(\alpha_A \otimes \alpha_B) \circ T = T \circ (\alpha_B \otimes \alpha_A)$. Then for any $a, a' \in A$ and $b \in B$,

$$\begin{aligned}
&(\mu_A \otimes \alpha_B)(id \otimes T)(T(b \otimes a) \otimes a') \\
=&\alpha_A^{-2}(a_{21}a'_{21}) \otimes \alpha_B^{-2}(b_{221})(S^{-1}\alpha_A^{-1}(a_1a'_1), \alpha_B^{-3}(b_{222}))(a'_{22}, \alpha_B^{-1}(b_{211}))(a_{22}, b_1) \\
=&\alpha_A^{-2}(a_{21}a'_{21}) \otimes \alpha_B^{-1}(b_{21})(S^{-1}\alpha_A^{-1}(a_1a'_1), \alpha_B^{-2}(b_{22}))(a_{22}a'_{22}, \alpha_B^{-1}(b_1)) \\
=&\alpha_A^{-2}(a_{21}a'_{21}) \otimes \alpha_B^{-1}(b_{12})(S^{-1}(a_1a'_1), b_2)(a_{22}a'_{22}, b_{11}) \\
=&T(\alpha_B(b) \otimes aa').
\end{aligned}$$

Similarly we have $T \circ (\mu_B \otimes \alpha_A) = (\alpha_A \otimes \mu_B) \circ (T \otimes id)(id \otimes T)$. Therefore T is a Hom-twisting map.

This finishes the proof. ■

Hence we have the Hom-associative algebra $A \bowtie B$ with the multiplication

$$(a \bowtie b)(a' \bowtie b') = (S^{-1}\alpha_A(a'_1), b_2)(a'_{22}, \alpha_B^{-1}(b_{11}))a\alpha_A^{-2}(a'_{21}) \bowtie \alpha_B^{-2}(b_{12})b'.$$

Theorem 3.4. If we define $\Delta : A \bowtie B \longrightarrow A \bowtie B \otimes A \bowtie B$ by

$$\Delta(a \bowtie b) = a_1 \bowtie b_2 \otimes a_2 \bowtie b_1,$$

the antipode by

$$S = T \circ \tau \circ (S_A \otimes S_B^{-1}).$$

Then $A \bowtie B$ is a Hom-Hopf algebra, which is called the Drinfeld double of A and B.

Remark 3.5. (1) Note that in the our construction, we don't need the Hom-Hopf algebra to be finite dimensional.

(2) In fact for any finite dimensional Hom-Hopf algebra (H, α), (H^{op}, H^*) is a dual pair if for any $h \in H$ and $f \in H^*$, $(a, f) = \langle f, a \rangle$. One can see that the Drinfeld double in this section is really a generalization of that in Corollary 2.16.

Proposition 3.6. By the multiplication, we can see that A and B could be imbedded into $A \bowtie B$ by $A \hookrightarrow A \bowtie B,\ a \mapsto a \bowtie 1_B$ and $B \hookrightarrow A \bowtie B,\ b \mapsto 1_A \bowtie b$, respectively. Moreover $a \bowtie b = (\alpha_A^{-1} \otimes \alpha_B^{-1})((a \bowtie 1_B)(1_A \bowtie b))$.

4. THE DRINFEL'D DOUBLE VERSUS THE HEISENBERG DOUBLE FOR HOM-HOPF ALGEBRAS

In this section, the relation between the Drinfel'd double and Heisenberg double is established.

Definition 4.1. Let (H, α) be a Hom-Hopf algebra. The linear map $\sigma : H \otimes H \longrightarrow k$ is called a *left Hom-2-cocycle* if the following conditions are satisfied

$$\sigma \circ (\alpha \otimes \alpha) = \sigma, \tag{4.1}$$

$$\sigma(l_1, k_1)\sigma(\alpha^2(h), l_2k_2) = \sigma(h_1, l_1)\sigma(h_2l_2, \alpha^2(k)), \tag{4.2}$$

for any $h, l, k \in H$.

Furthermore, σ is *normal* if $\sigma(1, h) = \sigma(h, 1) = \varepsilon(h)$.

Similarly if the condition (4.2) is replaced by

$$\sigma(\alpha^2(h), l_1k_1)\sigma(l_2, k_2) = \sigma(h_1l_1, \alpha^2(k))\sigma(h_2, l_2),$$

then σ is a *right Hom-2-cocycle*.

Example 4.2. (1) Recall from Corollary 2.15 that the Drinfeld double $D(H)$ of (H, α_H) is the space $H^{op} \otimes H^*$ with the multiplication

$$(h \otimes f)(k \otimes g) = \alpha_H^{-2}(k_{21})h \otimes [\alpha_H^{-3}(k_{22}) \rightharpoonup ((\alpha_H^*)^2(f) \leftharpoonup S\alpha_H^{-3}(k_1))]g,$$

for any $h, k \in H$ and $f, g \in H^*$.

Define $\sigma : D(H) \otimes D(H) \longrightarrow k$ by

$$\sigma(h \otimes f, k \otimes g) = \varepsilon(h)g(1)\langle f, \alpha(k)\rangle \tag{4.3}$$

Then it is not hard to verify that σ is a left Hom-2-cocycle on $D(H)$.

(2) In [8] Makhlouf and Panaite introduced another form of the Drinfeld double $\widetilde{D(A)}$, which is the space $(A^{op})^* \otimes A$ with the multiplication:

$$(f \otimes a)(g \otimes b) = f[(\alpha_A^{-3}(a_1) \rightharpoonup (\alpha_A^2(g))) \leftharpoonup S^{-1}\alpha_A^{-3}(a_{22})] \otimes \alpha_A^{-2}(a_{21})b,$$

for any $a, b \in A$ and $f, g \in A^*$.

Define $\eta : \widetilde{D(A)} \otimes \widetilde{D(A)} \longrightarrow k$ by

$$\eta(f \otimes a, g \otimes b) = \varepsilon(b)f(1)\langle g, \alpha(a)\rangle \tag{4.4}$$

Then η is a right Hom-2-cocycle on $\widetilde{D(A)}$.

Proposition 4.3. Let (H, α) be a Hom-Hopf algebra.

(1) If σ is a left Hom-2-cocycle, for any $h, k \in H$, define multiplication on H as follows

$$h \cdot_\sigma k = \sigma(h_1, k_1)\alpha^{-1}(h_2k_2),$$

then $(H, \cdot_\sigma, \alpha)$ is a Hom-associative algebra, called the left twist of H and denoted by ${}_\sigma H$.

(2) If σ is a right Hom-2-cocycle, define multiplication on H as follows

$$h \ {}_\sigma \cdot k = \alpha^{-1}(h_1k_1)\sigma(h_2, k_2),$$

then $(H, \cdot_\sigma, \alpha)$ is also a Hom-associative algebra, called the right twist of H and denoted by H_σ.

Proof. (1) First of all, for any $h, k, l \in H$, we have

$$1 \cdot_\sigma h = h \cdot_\sigma 1 = \alpha(h),$$

and

$$\alpha(h \cdot_\sigma k) = \sigma(h_1, k_1)\alpha(\alpha^{-1}(h_2k_2)) \overset{(4.1)}{=} \alpha(h) \cdot_\sigma \alpha(k).$$

Then

$$\begin{aligned}
\alpha(h) \cdot_\sigma (k \cdot_\sigma l) &= \alpha(h) \cdot_\sigma (\sigma(k_1, l_1)\alpha^{-1}(k_2l_2)) \\
&= \sigma(k_{11}, l_{11})\sigma(\alpha^2(h_1), k_{12}l_{12})h_2\alpha^{-1}(k_2l_2) \\
&\overset{(4.2)}{=} \sigma(h_{11}, k_{11})\sigma(h_{12}k_{12}, \alpha^2(l_1))\alpha^{-1}(h_2k_2)l_2 \\
&= (h \cdot_\sigma k) \cdot_\sigma \alpha(l).
\end{aligned}$$

(2) Similar to the proof of the part (1).

This finishes the proof. ■

Definition 4.4. The Heisenberg double $H(A)$ of a Hom-Hopf algebra (A, α_A) is the smash product $A\#A^*$ with respect to the left regular action of A^* on A, that is, it possesses the product as follows:

$$(a\#f)(b\#g) = a(f_1 \circ \alpha_A^2 \rightharpoonup \alpha_A^{-1}(b))\#(f_2 \circ \alpha_A)g,$$

for any $a, b \in A$ and $f, g \in A^*$.

We now have the following main result of this section.

Theorem 4.5. Let (A, α_A) be a Hom-Hopf algebra.

(1) The Heisenberg double $H(A^{op})$ of A^{op} is the left twist of the Drinfeld double $D(A)$ with the Hom-2-cocycle σ given by Eq.(4.3).

(2) The Heisenberg double $H(A^*)$ of A^* is the right twist of the Drinfeld double $\widetilde{D(A)}$ with the right Hom-2-cocycle η given by Eq.(4.4).

Proof. (1) We need to check that ${}_\sigma D(A)$ and $H(A^{op})$ share the same multiplication. Indeed for any $f, g \in A^*$ and $a, b \in A$,

$$\begin{aligned}
(a \otimes f) \cdot_\sigma (b \otimes g) &= \sigma(a_1 \otimes f_1, b_1 \otimes g_1)(\alpha^{-1} \otimes \alpha^*)((a_2 \otimes f_2)(b_2 \otimes g_2)) \\
&= \varepsilon(a_1)\alpha^{-3}(b_{221})\alpha^{-1}(a_2)\langle f_1, \alpha(b_1)\rangle \\
&\qquad \otimes [\alpha^{-4}(b_{222}) \rightharpoonup ((\alpha^3)^*(f_2) \leftharpoonup S\alpha^{-4}(b_{21}))]g \\
&= \alpha^{-1}(b_1)a \otimes \langle f, \alpha^{-1}(?b_2)\rangle g \\
&= (a\#f)(b\#g)
\end{aligned}$$

(2) We just compute the multiplication in $\widetilde{D(A)}$. For any $f, g \in A^*$ and $a, b \in A$,

$$\begin{aligned}
(f \otimes a) \cdot_\eta (g \otimes b) &= (\alpha^* \otimes \alpha^{-1})((f_1 \otimes a_1)(g_1 \otimes b_1))\eta(f_2 \otimes a_2, g_2 \otimes b_2) \\
&= f[(\alpha^{-4}(a_{11}) \rightharpoonup (\alpha^3)^*(g_2)) \leftharpoonup S^{-1}\alpha^{-4}(a_{122})] \\
&\qquad \otimes \alpha^{-3}(a_{121})b\langle f_1, \alpha(a_2)\rangle \\
&= f\langle g, \alpha^{-1}(?a_1)\rangle \otimes \alpha^{-1}(a_2)b,
\end{aligned}$$

which is exactly the multiplication in $H(A^*)$.

This completes the proof. ■

Example 4.6. Let (kG, ϕ) be the Hom-Hopf algebra in Example 2.17 (2). By Example 4.2 (1), we have a left Hom-2-cocycle σ on $D(kG) \otimes D(kG)$ given by

$$\sigma(g^i \otimes e_j, g^m \otimes e_k) = \delta_{k,0}\delta_{j,n-m},$$

for any $0 \leq i, j, m, k < n$. Then

$$\begin{aligned} & (g^i \otimes e_j) \cdot_\sigma (g^m \otimes e_k) \\ = \ & \sigma(g^{n-i} \otimes (e_j)_1, g^{n-m} \otimes (e_k)_1)(\phi^{-1} \otimes \phi^*)[(g^{n-i} \otimes (e_j)_2)(g^{n-m} \otimes (e_k)_2)] \\ = \ & (g^i \otimes e_{j+m})(g^m \otimes e_k) = g^{n-(i+m)} \otimes \delta_{j+m,k} e_{n-k} \\ = \ & (g^i \# e_j)(g^m \# e_k), \end{aligned}$$

just as shown in Theorem 4.5.

The following property is straightforward.

Proposition 4.7. Assume that σ is a left Hom-2-cocycle on the Hom-Hopf algebra (H, α), then the comultiplication Δ of H makes ${}_\sigma H$ into a right H-comodule Hom-algebra. Similar result holds for the algebra H_σ if σ is a right Hom-2-cocycle.

We now can apply Theorem 4.5 and Proposition 4.7 to obtain the following corollary:

Corollary 4.8. (1) The comultiplication of $D(A)$, considered as a map from $H(A^{op})$ to $H(A^{op}) \otimes D(A)$, makes $H(A^{op})$ into a right $D(A)$-comodule Hom-algebra.

(2) The comultiplication of $\widetilde{D(A)}$, considered as a map from $H(A^*)$ to $\widetilde{D(A)} \otimes H(A^*)$, makes $H(A^*)$ into a left $\widetilde{D(A)}$-comodule Hom-algebra.

ACKNOWLEDGEMENTS

This work was supported by the NSF of China (No. 11371088) and the NSF of Jiangsu Province (No. BK2012736).

References

[1] S. Caenepeel and I. Goyvaerts, Monoidal Hom-Hopf algebras, Comm. Alg., 39(2011), 2216–2240.

[2] V.G. Drinfeld, Quantumgroups, Proc. Internat. Congr. Math. Berkeley, 1(1986): 789–820.

[3] L. Delvauxa, A. Van Daele, The Drinfeld double versus the Heisenberg double for an algebraic quantum group, J. Algebra, 190(2004):59–84.

[4] L. Liu and B. Shen, Radford's biproducts and Yetter-Drinfel'd modules for monoidal Hom-Hopf algebras, J. Math. Phys. 55(2014), 031701.

[5] J. H. Lu, On the Drinfeld double and the Heisenberg double of a Hopf algebra, Duke Math. J., 74(1994): 763–776.

[6] S. Majid, Physics for Algebraists: Non-commutative and Non-cocommutative Hopf Algebras by a Bicrossproduct construction, J. Alg., 130(1990), 17–64.

[7] S. Majid, Foundations of quantum group theory, Cambridge university press, 1995.

[8] A. Makhlouf and F. Panaite, Hom-L-R-smash products, Hom-diagonal crossed products and the Drinfel'd double of a Hom-Hopf algebra, arXiv:1403.7077v1

[9] A. Makhlouf and S. D. Silvestrov, Hom-algebra structure, J. Gen. Lie Theory Appl. 2(2008), 52–64.

[10] A. Makhlouf and S. D. Silvestrov, Hom-algebras and Hom-coalgebras, J. Alg. Appl., 9(2010), 553–589.

[11] S. H. Wang, Turaev Group Coalgebras and Twisted Drinfel'd Double, Indiana University Mathematics Journal, 58(2009): 1395–1417.

[12] D. Yau, Hom-quantum groups: I. Quasi-triangular Hom-bialgebras, J. Phys. A: Math. Theor., 45(2012), 065203.

[13] M. M. You, S. H. Wang, Constructing New Braided T-categories over Monoidal Hom-Hopf Algebras, J. Math. Phys.55(2014), 111701.